184-34388

NASA Technical Memorandum

NASA TM-86458

ON LONG-TERM PERIODICITIES IN THE SUNSPOT RECORD

By Robert M. Wilson Space Science Laboratory

July 1984



George C. Marshall Space Flight Center

	TECHNIČ	CAL REPORT STANDARD TITLE PAGE
1. REPORT NO.	2. GOVERNMENT ACCESSION NO.	3. RECIPIENT'S CATALOG NO.
NASA TM-86458		
4. TITLE AND SUBTITLE		5. REPORT DATE
	<i>(</i>	July 1984
On Long-Term Periodicities i	in the Sunspot Record	6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S)	and a second section of the second	8. PERFORMING ORGANIZATION REPORT #
Robert M. Wilson		WORK THAT NO
9. PERFORMING ORGANIZATION NAME AND A	ADDRESS	10. WORK UNIT, NO.
		CONTRACT OF CRANT NO
George C. Marshall Space Fl		11. CONTRACT OR GRANT NO.
Marshall Space Flight Center	Alabama 35812	13. TYPE OF REPORT & PERIOD COVERED
10 COONCODING ACENCY NAME AND ADDRESS		13. TIPE OF REPORT & PERIOD COVERED
12. SPONSORING AGENCY NAME AND ADDRES		market al Manager de la constant
National Assessment on a Con	a a A ductod obrobiom	Technical Memorandum
National Aeronautics and Spa	ace Administration	14. SPONSORING AGENCY CODE
Washington, D.C. 20546		,
15. SUPPLEMENTARY NOTES		ang pangkang ang ang ang ang ang ang ang ang ang
Prepared by Space Science	Laboratory, Science and Engin	neering
16. ABSTRACT		
an 11-year average period en highest quality and consider present. On the basis of configuration of sine curves to approximate smoothed sunspot number at sinusoidal function, having periodicity, yields computed	been systematically maintained exists, since about 1850. Thused to be the most reliable is sycles 8 through 20 , we have use the observed $\overline{R}_{\text{MAX}}$ values cycle maximum). We find that an 11-cycle and a 2-cycle variance $\overline{R}_{\text{MAX}}$ values which fit, reasonable.	s, the sunspot record of that of cycle 8 through the used various combinations (where \overline{R}_{MAX} is the at a three-component iation on a 90-cycle onably well, observed
	n sunspot cycles. Extrapolation	
forward in time allows us to	project values of \overline{R}_{MAX} for c	cycles 21 and 22. For cycle
of 164.5 and to that predict function projects a value of have been applied to cycle a period cycle (cycle duration)	value of 157.3, very close to ed earlier by Sargent [64] - about 107. Linear regression 22, yielding the result that it > 132 months). A major feat echniques used to estimate the	154. For cycle 22, the as identified in Wilson [44] will probably be a longure of this report is an
		•

17. KEY WORDS Sun, Sunspots, Solar Cycle, Sunspot Periodicities		Unclassified	il_	
19. SECURITY CLASSIF. (of this report)	20. SECURITY CLAS	SIF. (of this page)	21. NO. OF PAGES	22. PRICE
Unclassified	Unclass	sified	48	NTIS

ACKNOWLEDGMENTS

The author is grateful to Mr. Jesse B. Smith, Jr. (NOAA/SEL) for reading the manuscript and offering helpful comments.

TABLE OF CONTENTS

		Page
I.	INTRODUCTION	1
II.	APPROACH	2
III.	RESULTS AND DISCUSSION	4
	A. Linear-Regression Fits (APPROACH 1) B. 50-Cycle Fits (APPROACH 2) C. 90-Cycle Fits (APPROACH 3) D. Pre-Cycle 8 Fits E. Cycles Beyond SCN 20	9 11 14 18 23
IV.	CONCLUSIONS	26
APPE	NDIX — LINEAR REGRESSIONS BASED ON \overline{R}_{MAX}	27
REFÉ	RENCES	31

LIST OF ILLUSTRATIONS

Figure	Title	Page
1.	$\overline{R}_{ ext{MAX}}$ values versus SCN (cycles 8 through 20)	5
2.	Statistical spread versus SCN (cycles 8 through 20)	8
3.	LR + 8 FIT	9
4.	LR + 11 FIT	10
5.	LR + 11 + 8 FIT	11
6.	LR + 11 + 2 FIT	12
7.	50 + 8 FIT	12
8.	50 + 11 FIT	13
9.	50 + 11 + 8 FIT	14
10.	50 + 8 + 2 FIT	15
11.	50 + 11 + 2 FIT	15
12.	90 + 8 FIT	16
13.	90 + 11 FIT	17
14.	90 + 8 + 2 FIT	17
15.	90 + 11 + 2 FIT	18
16.	Summary chart: Computed \overline{R}_{MAX} values for SCN -64 to SCN 30	19
17.	R(A) versus \overline{R}_{MAX}	21
18.	$ ext{SLOPE}_{ ext{ASC}}, \ \overline{ ext{R}}_{ ext{CHM}}, \ \overline{ ext{R}}_{ ext{MEAN}} \ ext{and} \ \overline{ ext{R}}_{ ext{MIN}} \ ext{versus} \ \overline{ ext{R}}_{ ext{MAX}}$	27
19.	MIN-MIN PERIOD, D_{CHM} and ASC_{CHM} versus \overline{R}_{MAX}	28
20.	DES and ASC versus $\overline{R}_{ ext{MAX}}$	29
21.	SLOPE _{DES} versus SLOPE _{ASC}	30
22.	s _{MEAN} versus \bar{R}_{MEAN}	30

LIST OF TABLES

Table	Title	Page
1.	Selected Parametric Values for Cycles 8 Through 20	3
2.	Angular Values (Phases) for Selected Periodic Cycles as a Function of SCN	6
3.	Selected Parametric Values for Cycles 1 Through 7	20
4.	Comparison of Reported and Computer R _{MAX} Values for the Maunder Minimum	22
5.	\overline{R}_{MAX} (COMP) Values for SCN 21	23
6.	Selected Parametric Values for SCN 21	24
7.	R _{MAX} (COMP) Values for SCN 22	25
8.	Selected Parametric Values for SCN 22	25

LIST OF SELECTED ABBREVIATIONS

Meaning Abbreviation

Ascent cycle duration in months ASC

Time in months from $\overline{R}_{\mbox{\footnotesize{MIN}}}$ to $\overline{R}_{\mbox{\footnotesize{MAX}}};$ CHM means "corrected half maximum" ASC_{CHM}

Time in months when $\overline{R}_{13} \stackrel{>}{=} \overline{R}_{CHM}$ D_{CHM}

DES Descent cycle duration in months

Linear Regression LR

Minimum-to-minimum cycle duration; ASC + DES MIN-MIN PERIOD

N Number

Smoothed sunspot number; monthly moving average of sunspot \overline{R}_{13}

R(A) Annual mean sunspot number

 $(\overline{R}_{MAX} + \overline{R}_{MIN})/2$ $\overline{R}_{\text{CHM}}$

 $\overline{R}_{\text{MAX}}$ \overline{R}_{13} value at cycle maximum

 $\overline{R}_{MAX}(COMP)$ Computed \overline{R}_{MAX}

Observed \overline{R}_{MAX} $\overline{R}_{MAX}(OBS)$

 \overline{R}_{MEAN} Mean \overline{R}_{13} value for entire cycle

 \overline{R}_{13} value at cycle minimum \overline{R}_{MIN}

Standard deviation S

Standard deviation associated with \overline{R}_{MEAN} SMEAN

 $SLOPE_{ASC}$ $(\overline{R}_{\text{MAX}}$ - $\overline{R}_{\text{MIN}})/\text{ASC},$ where all parameters are for same cycle

 $SLOPE_{DES}$ $(\overline{R}_{MIN}$ - $\overline{R}_{MAX})$ /DES, where \overline{R}_{MAX} and DES are for a particular

cycle and \overline{R}_{MIN} is for the succeeding cycle

TECHNICAL MEMORANDUM

ON LONG-TERM PERIODICITIES IN THE SUNSPOT RECORD

I. INTRODUCTION

In recent years, substantial interest has been given to the sunspot record, especially its historical aspects. Much of this work sought to associate possible climatic excursions with variations in sunspot number. For example, Eddy [1-8] has done considerable research on the historical record of sunspot data and solar activity (also [9-43, 162-167, 172-183]). His studies, based primarily on naked-eye and early telescopic sunspot observations, fossil radiocarbon levels in tree rings, and auroral reports, have suggested possibly strong climatic associations in the Sun-Earth record; in particular, he has found evidence for the occurrence of prolonged "sunspot minima," e.g., the Spörer Minimum (A.D. 1400-1510) and Maunder Minimum (A.D. 1645-1715), and for the occurrence of a prolonged "sunspot maximum," i.e., the so-called Medieval Maximum or Grand Maximum (ca., A.D. 1120-1280). In addition, he has given tutorial remarks concerning the sunspot record post-1850 and pre-1850; i.e., he notes that the highest or best-quality sunspot data are those dated from about 1850 to the present, since it was about 1850 when solar observers first began to record telescopic sunspot observations in a systematic fashion, having just become aware of the existence of the cyclic nature of sunspots (Wilson [44] and Wilson et al. [45]). Sunspot observations prior to about 1850 are viewed as being somewhat less reliable, or at least somewhat different, as a simple comparison of selected cycle parameters for pre-1850 to those post-1850 will show. (Sonett [36] has suggested that there may be "bookkeeping" type errors in the compilations associated with these early sunspot cycles; also it must not be forgotten that daily sunspot records were not maintained on a continuous basis until January 1849, Waldmeier [170].) Cycles have been arbitrarily assigned numbers back in time into the 1700's (and before), with sunspot cycle number (SCN) 1 having a maximum in June 1761; the present cycle, now in decline, is denoted SCN 21.

Another aspect of this work has been to estimate or "predict" sunspot numbers as a function of time for future epochs, based on modeling or "post-dicting" of past This aspect usually incorporates various time series techniques which sometimes make use of power-spectrum analysis results. Power-spectrum analyses suggest the existence of several possible short-term and long-term periodicities in the solar cycles; e.g., several authors have reported the presence of 5-, 8-, 10-, 22-, 90-, and 180-year periods, as well as the well-known 11-year average period cycle duration in the sunspot record. Some authors have hypothesized that these periods may be due to planetary motion "tidal" effects, while others argue that sunspot-like cycles can be simulated with random noise. It must be pointed out, however, that some skepticism remains regarding the authenticity of most of these short-term and longterm periodicities, and, following Wallenhorst [62], it is important to note that the 11-year periodicity is the only statistically significant one present in the sunspot We should recall that sunspots have been routinely and systematically monitored for a mere 130 years or so; thus, few or no cycles of periods substantially longer than 11 years have been observed. Wilson [44], Wallenhorst [46], Newkirk and Frazier [47], Bray and Loughhead [48], Noyes [49], and Meadows [50] have given additional comments regarding sunspot numbers, their history, measurement, and use.

A number of references [51-161, 170-183] are included which allows the reader to follow the past 80 years growth in our understanding of sunspot cyclicity and to see the methods employed for estimating contemporary values of sunspot number.

The purpose of this report is threefold: (1) to compare observed sunspot numbers at cycle maximum of contemporary cycles (i.e., cycles 8 through 20) with similar computed values, based on the assumption that sunspots are periodic in nature (i.e., sinusoidal); (2) to extrapolate the preferred, empirical periodic function "backwards" in time to qualitatively compare computed values with values estimated by others for such epochs as the Maunder Minimum; and (3) to extrapolate the preferred periodic function "forwards" in time to estimate smoothed sunspot number at sunspot maximum (denoted, \overline{R}_{MAX}) for cycles 21, 22, and beyond. In addition, using linear relationships identified in Wilson [44], a number of cycle-related parametric values will be estimated and compared with observed values (when they are known) for cycle 21 and will be projected for cycle 22. In Section II, the approach used in this study is described; in Section III, the results and discussion of the findings are presented; and in Section IV, the conclusions are stated. This report is a continuation of studies investigating solar activity and its estimate (e.g., Wilson [44, 102], Wilson et al. [45, 100], and Teuber et al. [107]).

II. APPROACH

Since Eddy [4] (and before him McNish and Lincoln [123]) has cautioned that cycles observed prior to about 1850, corresponding to SCN 9, may not be reliably known, many solar statisticians today limit their remarks concerning cycle-related parameters, such as ascent period (dubbed ASC, the time in months measured from \overline{R}_{MIN} occurrence to \overline{R}_{MAX} occurrence; \overline{R}_{MIN} is the minimum smoothed sunspot number or \overline{R}_{13} value marking the beginning of a cycle and \overline{R}_{MAX} is the maximum \overline{R}_{13} value in the same cycle), descent period (dubbed DES, the time in months from \overline{R}_{MAX} occurrence to subsequent cycle \overline{R}_{MIN} occurrence), cycle duration (denoted MIN-MIN PERIOD or the cycle minimum-to-minimum period, numerically equal to ASC + DES), and others to cycles 8 through 20. Table 1 lists values for the cycle-related parameters for cycles 8 through 20. It also gives mean and "uncorrected" standard deviation for each parameter. Definitions of these parameters are given in the list of selected abbreviations found at the beginning of this report.

Using \overline{R}_{MAX} values for cycles 8 through 20, this report compares a linear-regression approach and modifications to it (APPROACH 1) with two cyclic approaches and modifications to them (APPROACH 2 and APPROACH 3). The two cyclic approaches presume a 50-cycle or approximately 550-year periodicity and a 90-cycle or 990-year periodicity in sunspot number, respectively. The reasoning behind the selection of these two periods will become evident in the following.

If restricted to cycles 8 through 20, it is observed that the mean value for $\overline{R}_{\rm MAX}$ is 116.2 with a 1-sigma confidence value (or standard deviation, s) of 36.7. (For small sample size, N, when N < 50, the "corrected" 1-sigma value is equal to $\sqrt{N/(N-1)}$ x s, or 38.2.) Thus, any given cycle can "crudely" be predicted to have, assuming a 1-sigma confidence with no apparent trend in the data, an $\overline{R}_{\rm MAX}$

TABLE 1. SELECTED VALUES FOR CYCLES 8 THROUGH 20

sc	N R _{MAX}	R _{MIN}	R _{MEAN}	SMEAN	R _{CHM}	ASCCHM	D _{CHM}	ASC	DES	MIN-MIN PERIOD	SLOPEASC	SLOPEDES
8	146.9	7.3	66.9	43.4	77.1	23	53	40	76	116	3.490	-1.795
9	132.0	10.5	55.9	30.4	71.3	44	39	55	94	149	2.209	-1.370
10	97.9	3.2	48.2	29.6	50.6	29	57	50	85	135	1.894	-1.091
11	140.5	5.2	52.9	43.8	72.9	28	46	41	100	141	3.300	-1.383
12	74.6	2.2	33.8	23.8	38.4	22	63	60	74	134	1.207	-0.941
13	87.9	5.0	38.5	28.1	46.5	20	54	47	96	143	1.764	-0.888
14	64.2	2.7	32.1	22.3	33.5	23	72	49	89	138	1.255	-0.704
15	105.4	1.5	44.0	30.5	53.5	27	50	49	71	120	2.120	-1.406
16	78.1	5.6	40.3	25.6	41.9	24	59	57	65	122	1.272	-1.148
17	119.2	3.5	57.2	36.9	61.4	29	57	43	82	125	2.691	-1.360
18	151.8	7.7	74.0	50.5	79.8	27	51	39	83	122	3.695	-1.788
19	201.3	3.4	90.5	67.6	102.4	23	54	47	79	126	4.211	-2.427
20	110.6	9.6	59.9	34.8	60.1	23	74	49	91	140	2.061	-1.081
MEAN ₈₋₂₀	116.2	5.2	53.4	35.9	60.7	26.3	56.1	48.2	83.5	131.6	2.398	-1.337
^{\$} 8–20	36.7	2.7	16.2	_	18.8	5.8	9.2	6.2	10.0	10.0	0.958	0.441

value equal to approximately 116.2 \pm 36.7. Statistically, correct prediction would be expected about 68 percent of the time. A 2-sigma confidence, or $\overline{R}_{MAX} = 116.2 \pm 73.4$, should result in being correct about 95 percent of the time. Therefore, it would be extremely useful to deduce a relationship which gives better estimates for \overline{R}_{MAX} with reduced 1-sigma spreads. In APPROACH 1, \overline{R}_{MAX} values are "detrended" for cycles 8 through 20 by means of standard linear-regression time series analysis (e.g., Downie and Heath [168] and Longley-Cook [169]).

Examination of \overline{R}_{MAX} values for cycles 8 through 20 clearly shows a downward trend between cycles 8 through about 14 and an upward trend from cycle 14 through at least cycle 19. The entire period spanning cycles 8 through 20 shows an upward trend. Based purely on the deduced regression equation, \overline{R}_{MAX} can be "better" predicted since s has been reduced to 33.2. Various modifications will be made on this linear-regression fit (denoted LR) to further reduce s. These modifications include assuming the presence of an 8-cycle periodicity or an 11-cycle periodicity, an 11-cycle periodicity with an imbedded 8-cycle periodicity, and an 11-cycle periodicity with an imbedded 2-cycle periodicity. The 8-cycle periodicity was selected because it is equivalent to the so-called "Gleissberg" 90-year period often said to be imbedded in the sunspot data [55]. The 11-cycle periodicity is the "best-fit" periodicity for the given \overline{R}_{MAX} values (cycles 8 through 20). (The term "best-fit" is used extensively in this report. By this we mean that the fit to the observed values of \overline{R}_{MAX} has the least sum of the squares of the residuals; see Section III.) The 2-cycle periodicity was selected since throughout much of the period spanning cycles 8 through 20, there is an apparent "up-down-up" signal in the data.

In APPROACH 2, \overline{R}_{MAX} values are fitted for eyeles 8 through 20 using a sine curve with a period equal to 50 cycles. This period was chosen based on the assumption that the Maunder Minimum actually occurred (the middle of the minimum corresponds to SCN -6) and that the sine curve is at phase 270 deg at that point and at phase 90 deg at SCN 19; hence, the half-period of the sine curve is 25-cycles, yielding the period to be 50 cycles. Such an analysis gives a 1-sigma value of 37.0 and yields a line that is very close in value to the linear-regression line. Modifications to this approach are as above in APPROACH 1, using various combinations of an 11-cycle, 8-cycle, and/or 2-cycle periodicity imbedded in the 50-cycle fit.

In APPROACH 3, the LR fit is approximated by a sine curve of period equal to 90 cycles, such that the curve is at phase 90 deg during the Grand Maximum (at SCN -53) and at phase 270 deg during the Maunder Minimum (at SCN -8). Such an analysis gives a 1-sigma value of 36.1 and yields a line very close to that of the regression line. Modifications to this approach follow that above for APPROACH 1 and APPROACH 2, using various combinations of an 11-cycle, 8-cycle, and/or 2-cycle periodicity imbedded in the 90-cycle fit.

III. RESULTS AND DISCUSSION

Following the approaches identified in the previous section, several fits to the observed \overline{R}_{MAX} values for cycles 8 through 20 are obtained. These basic, first-order fits are shown in Figure 1. The observed values are plotted as darkened circles and the computed values as darkened triangles. The lines shown are: mean fit, linear-regression fit, 39-cycle fit, and 90-cycle fit: Equations are given below and in Figure 1 for these fits:

Mean Fit

$$\overline{R}_{MAX} = 116.2 \tag{1}$$

Linear-Regression Fit

$$\overline{R}_{MAX} = 88.1 + 2.0 X \tag{2}$$

50-Cycle Fit

$$\bar{R}_{MAX} = 100.7 + 18.7 \sin \theta_1(X)$$
 (3)

90-Cycle Fit

$$\overline{R}_{MAX} = 116.0 + 24.7 \sin \theta_5(X)$$
 (4)

In equation (2), X is the sunspot cycle number (SCN). In equations (3) and (4), $\theta_1(X)$ is the angle in degrees for a particular X assuming a 50-cycle periodicity, and




Figure 1. \overline{R}_{MAX} values versus SCN (cycles 8 through 20).

ř,

 $\theta_5(X)$ is the angle in degrees for a particular X assuming a 90-cycle periodicity, respectively. These angular values are identified in Table 2. The standard deviations for each of these fits are 36.7, 36.1, 37.0, and 36.1, respectively. Equation (3) is of the form $y=a_0+a_1\sin\theta$, where $a_0=100.7$ and $a_1=18.7$. The constant a_0 was chosen to be 100.7 on the basis that SCN 19 has the highest known \overline{R}_{MAX} value (= 201.3) and that the lowest possible value for any cycle is 0; thus, a_0 is the rounded-off half-way point between these two extremes. The constant a_1 is the "best-fit" value based on the given a_0 and the values of $\theta_1(X)$ as shown in Table 2 for cycles 8 through 20. The "best-fit" a_1 was deduced by arbitrarily selecting three values of a_1 and computing \overline{R}_{MAX} values for cycles 8 through 20 with the given a_0 , summing the squares of the residuals (where a residual is the difference between the observed and computed value of \overline{R}_{MAX} for a cycle), and fitting these sums of squares of the residuals versus a_1 which yields the least value for the sums of the squares of the residuals.

TABLE 2. ANGULAR VALUES (PHASES) FOR SELECTED PERIODIC CYCLES AS A FUNCTION OF SCN

		50-CYCLE	11-CYCLE	8-CYCLE	2-CYCLE	90-CYCLE	
		×	ξ	ĕ	×	∑	
	01/01/5 1141//14/11	ပု	ပု	٩	ပု	ပို	
	CYCLE MAXIMUM APPROXIMATE	20	=	œ	6	8	
SCN	YEAR DATE	θ_1X	θ ₂ X	θ_3X	θ ₄ X	θ ₅ X	NOTES
-60	1088	241.2	24.5	135	270	62	
-59	1099	248.4	57.3	180	90	66	
-58	1110	255.6	90.0	225	270	70	
-57	1121	262.8	122.7	270	90	74	(APPROXIMATE)
-56	1132	270.0	155.5	315	270	78	1
-55	1143	277.2	188.2	360	90	82	
-54	1154	284.4	220.9	45	270	86	
53	1165	291.6	253.6	90	90	90	
-52	1176	298.8	286.4	135	270	94	GRAND MAXIMUM
-51	1187	306.0	319.1	180	90	98	
50	1198	313.2	351.8	225	270	102	
-49	1209	320.4	24.5	270	90	106	
-48	1220	327.6	57.3	315	270	110	
47	1231	334.8	90.0	360	90	114	
-46	1242	342.0	122.7	45	270	118	:
-45	1253	349.2	155.5	.90	90	122	
-44	1264	356.4	188.2	135	270	126	1
-43	1275	3.6	220.9	180	90	130	(45550)
-42	1286	10.8	253.6	225	270	134	(APPROXIMATE)
-41	1297	18.0	286.4	270	90	138	
-40	1308	25.2	319.1	315	270	142	
39	1319	32.4	351.8	360	90	146	
-38 -37	1330 1341	39.6	24.5	45	270	150	
-37 -36		46.8	57.3	90	90	154	
-35 -35	1352	54.0	90.0	135	270	158	
34	1363	61.2	122.7	180	90	162	
34 33	1374	68.4	155.5	225	270	166	
-32	1385 1396	75.6	188.2	270	90	170	
-32 -31	1407	82.8 90.0	220.9 253.6	315	270	174	
30	1418	97.2	286.4	360 45	90	178 182	(APPROXIMATE)
-29	1429	104.4	319.1	90	270 90	186	T (ALL HOXIMALE)
-28	1440	111.6	351.8	135	270	190	Ī
-27	1451	118.8	24.5	180	90	194	
-26	1462	126.0	57.3	225	270	198	SPÖRER
-25	1473	133.2	90.0	270	90	202	MINIMUM
-24	1484	140.4	127.7	315	270	202	I I I I I I I I I I I I I I I I I I I
-23	1495	147.6	155.5	360	90	210	
-22	1506	154.8	188.2	45	270	214	
-21	1517	162.0	220.9	90	90	218	(APPROXIMATE)
-20	1528	169.2	253.6	135	270	222	
-19	1539	176.4	286.4	180	90	226	
-18	1550	183.6	319.1	225	270	230	
-17	1561	190.8	351.8	270	90	234	
-16	1572	198.0	24.5	315	270	238	
,,,	1072	130.0	24.0	3 10	2/0	230	

TABLE 2. (Concluded)

	CYCLE MAXIMUM APPROXIMATE	50-CYCLE	11-CYCLE	8-CYCLE	2-CYCLE	90-CYCLE	
SCN	YEAR DATE	$\theta_1 X$	$\theta_2 X$	θ_3X	$\theta_{4}X$	$\theta_{5}\mathbf{X}$	NOTES
<u></u>	1583	205.2	57.3	360	90	242	
-14	1594	212.4	90.0	45	270	246	
-13	1605	219.6	122.7	90	90	250	
-12	1616	226.8	155.5	135	270	254	
-11	1627	234.0	188.2	180	90	258	(40000)(114.75)
-10	1638	241.2	220.9	225	270	262	(APPROXIMATE)
- 9	1649	248.4	253.6	270	90	266	Ţ
- 8	1660	255.6	286.4	315	270	270	MAUNDER
- 7	1671	262.8 270.0	319.1 351.8	360 45	90 270	274 278	MINIMUM
- 6 - 5	1682 1693	277.2	24.5	90	90	282	***************************************
– 5 – 4	1704	284.4	57.3	135	270	286	1
- 3	1715	291.6	90.0	180	90	290	(APPROXIMATE)
- 3 - 2	1726	298.8	122.7	225	270	294	
- ī	1737	306.0	155.5	270	90	298	
Ó	1748	313.2	188.2	315	270	302	
1	1759	320.4	220.9	360	90	306	
2	1770	327.6	253.6	45	270	310	
.3	1781	334.8	286.4	90	90	314	
4	1792	342.0	319.1	135	270	318	
5	1803	349.2	351.8	180	90	322	
6	1814	356.4	24.5	225	270	326	
7	1825	3.6	57.3	270	90	330	
8	1836	10.8	90.0	315	270	334	BEST-QUALITY
9	1847	18.0	122.7	360	90	338	SUNSPOT DATA
10	1858	25.2	155.5	45	270	342 346	JUNSIOT DATA
11	1869	32.4 39.6	188.2	90 135	90 270	350	
12 13	1880 1891	39.6 46.8	220.9 253.6	180	90	354	
14	1902	54.0	286.4	225	270	358	
15	1913	61.2	319.1	270	90	2	
16	1924	68.4	351.8	315	270	6	
17	1935	75.6	24.5	360	90	10	
18	1946	82.8	57.3	45	270	14	
19	1957	90.0	90.0	90	90	18	
20	1968	97.2	122.7	135	270	22	
21	1979	104.4	155.5	180	90	26	
22	1990	111.6	188.2	225	270	30	
23	2001	118.8	220.9	270	90	34	
24	2012	126.0	253.6	315	270	38	
25	2023	133.2	286.4	360	90	42	
26	2034	140.4	319.1	45	270	46 50	
27	2045	147.6 154.8	351.8 24.5	90 135	90 270	54	
28 29	2056 2067	162.0	24.5 57.3	180	90	58 58	
30	2078	169.2	90.0	225	270	62	

Similarly, equation (4) is of the form $y = a_0 + a_1 \sin \theta$, where $a_0 = 116$ and $a_1 = 24.7$. The constant a_0 was selected to be approximately the value that the linear-regression equation [eq. (2)] yields for SCN 14, the mid-point of the data set. As above, a_1 is the "best-fit" for the given a_0 and the values of $\theta_5(X)$ shown in Table 2, using the least-squares method.

In Figure 2, the residuals are plotted in units of standard deviation for each of the fits shown in Figure 1. For convenience, the figure is arbitrarily divided into several smaller regions. The middle region of each plot is designated "average" since approximately 50 percent, on average, of the \overline{R}_{MAX} values are expected to lie within its bounds; the region just above (and below) "average" is called "above (below)




Figure 2. Statistical spread versus SCN (cycles 8 through 20).

-2.0S

-ANOMALOUS LOW

(<5%)

–2.0_S

- ANOMALOUS LOW

(<5%)


average and approximately 20 percent, on average, of the cycles are expected to fall within its bounds. The top-most (bottom-most) region is called "anomalous high (low)" and represents the approximately upper (lower) 5 percent, on average, of the $\overline{R}_{\rm MAX}$ values which are greater than 1.645 standard deviations above (below) the computed value for $\overline{R}_{\rm MAX}$. Based on these single-component fits, SCN 19 is observed to probably be an anomalously high $\overline{R}_{\rm MAX}$ cycle.

In subsections A, B, and C which follow, the effects of selected modifications to the aforementioned basic curve-fits (linear regression, 50-cycle, and 90-cycle fits) are examined. In subsection D, computed values of $\overline{R}_{\rm MAX}$ are compared using the preferred empirical fit with values reported for earlier epochs (SCN 1 through 7 and the Maunder Minimum). (The reader is reminded that phase for the empirical fit is based, in part, on the occurrence of the Maunder Minimum; however, amplitude for the fit is based only on cycles 8 through 20.) In Subsection E, the empirical fit is used to infer values of $\overline{R}_{\rm MAX}$ for future epochs (SCN 21, 22, and beyond).

A. Linear-Regression Fits (APPROACH 1)

Following the first modification in APPROACH 1, Figure 3 shows the linear-regression fit with an 8-cycle variation. The equation for this fit is:

$$\overline{R}_{MAX} = 88.1 + 2.0 X + 19.8 \sin \theta_3(X)$$
 , (5)






Figure 3. LR + 8 FIT.

where X is again SCN and $\theta_3(X)$ is found in Table 2. The constant 19.8 is the "best-fit" for the values given in the table and above. This fit only slightly reduces the statistical spread (s = 33.4 for the LR + 8 FIT as compared to s = 36.1 for the LR FIT alone); so, if indeed there is an 8-cycle periodicity in the \overline{R}_{MAX} record for cycles 8 through 20, it does not appear to contribute much to the variation in \overline{R}_{MAX} value with time. Based on this two-component fit, both cycles 8 and 19 are observed to probably be anomalously high.


A better two-component fit is plotted in Figure 4 which shows the linear-regression fit with an 11-cycle variation. Its equation is:

$$\overline{R}_{MAX} = 88.1 + 2.0 X + 30.7 \sin \theta_2(X)$$
 (6)

As before, X is SCN and $\theta_2(X)$ can be found in Table 2. The constant 30.7 is the "best-fit" for the given table values and constants used in equation (6). This fit reduces the statistical spread by about 30 percent (s = 25.2 for the LR + 11 FIT). Based on this fit, cycle 19 is anomalously high and cycle 20 is anomalously low.

Figure 5 shows a three-component fit based on the linear-regression fit with an 11-cycle and 8-cycle variation. Its equation is:

$$\overline{R}_{MAX} = 88.1 + 2.0 X + 30.7 \sin \theta_2(X) + 11.4 \sin \theta_3(X) .$$
 (7)







Figure 4. LR + 11 FIT.






Figure 5. LR + 11 + 8 FIT.

Again, this equation only slightly reduces the statistical spread as compared to equation (6) (s=24.9 for the LR + 11 + 8 FIT). Based on this fit, cycle 20 is anomalously low.

The final fit following APPROACH 1 is shown in Figure 6. It is a three-component fit based on the linear-regression fit with an 11-cycle and 2-cycle variation. Its equation is:


$$\overline{R}_{MAX} = 88.1 + 2.0 X + 30.7 \sin \theta_2(X) + 15.8 \sin \theta_4(X)$$
 (8)

This fit reduces the original statistical spread from s=36.1 to s=19.7, a 45-percent reduction. (The variance has been reduced by 70 percent.) No cycles are observed to be anomalously low or high based on this fit, although several are in the above average or below average categories.

B. 50-Cycle Fits (APPROACH 2)

In APPROACH 2, the baseline fit is a 50-cycle periodicity. Modifications to it will be made as in APPROACH 1. Figure 7 depicts an 8-cycle variation on the 50-cycle fit. Its equation is:

$$\bar{R}_{MAX} = 100.7 + 18.7 \sin \theta_1(X) + 21 \sin \theta_3(X)$$
 (9)







Figure 6. LR + 11 + 2 FIT.







Figure 7. 50 + 8 FIT.

The statistical spread about this line is only slightly better than the 50-cycle fit alone (s = 33.8 for the 50 + 8 FIT). Cycles 8 and 19 both appear as anomalously high cycles.

In Figure 8, an 11-cycle variation on the 50-cycle fit is plotted. As in APPROACH 1, a much better fit is suggested. Its equation is:

$$\overline{R}_{MAX} = 100.7 + 18.7 \sin \theta_1(X) + 33.2 \sin \theta_2(X)$$
 (10)

Its statistical spread is s = 24.5, about 34 percent more restrictive than the 50-cycle fit alone. Cycle 19 is the only cycle that appears to be anomalously high.







Figure 8. 50 + 11 FIT.

In Figure 9, an 11-cycle and an 8-cycle variation have been placed on the 50-cycle fit. Its equation is:

$$\overline{R}_{MAX} = 100.7 + 18.7 \sin \theta_1(X) + 33.2 \sin \theta_2(X) + 8.7 \sin \theta_3(X)$$
 (11)

As before, the presence of an 8-cycle periodicity has little effect upon the reduction of the statistical spread (s = 23.8 for the 50 + 11 + 8 FIT). Cycles 19 and 20 appear to be anomalously high and low, respectively.






Figure 9. 50 + 11 + 8 FIT.

In Figure 10, an 8-cycle and 2-cycle variation has been included with the 50-cycle periodicity. Its equation is:

$$\overline{R}_{MAX} = 100.7 + 18.7 \sin \theta_1(X) + 21.9 \sin \theta_3(X) + 12.8 \sin \theta_4(X)$$
 (12)

This fit is worse than the preceding fit, having s = 31.2. Cycle 8 appears to be an anomalously high cycle.


A much better fit is that shown in Figure 11 where an 11-cycle and a 2-cycle variation has been added to the baseline 50-cycle fit. Its equation is:

$$\overline{R}_{MAX} = 100.7 + 18.7 \sin \theta_1(X) + 33.2 \sin \theta_2(X) + 15.4 \sin \theta_4(X)$$
 (13)

Its statistical spread is 49 percent smaller than that deduced for the baseline (s = 19.0 for the 50 + 11 + 2 FIT; the variance has been reduced by 74 percent). Only cycle 19 appears to be anomalously high.

C. 90-Cycle Fits (APPROACH 3)

In this subsection modifications to a 90-cycle periodicity, which approximates the linear-regression equation used in APPROACH 1, are examined. Figure 12 shows an 8-cycle variation on the 90-cycle fit. Its equation is:







Figure 10. 50 + 8 + 2 FIT.







Figure 11. 50 + 11 + 2 FIT.






Figure 12. 90 + 8 FIT.

$$\overline{R}_{MAX} = 116.0 + 24.7 \sin \theta_5(X) + 20.3 \sin \theta_3(X)$$
 (14)

Its spread is 33.1, only slightly reduced from the 90-cycle baseline spread of 36.1. Cycles 8 and 19 appear to be anomalously high.

A better fit is obtained using an 11-cycle variation on the 90-cycle periodicity, shown in Figure 13. Its equation is:


$$\overline{R}_{MAX} = 116.0 + 24.7 \sin \theta_5(X) + 35.4 \sin \theta_2(X)$$
 (15)

Its 1-sigma spread is 24.6 units. Cycles 19 and 20 appear to be anomalously high and low, respectively.

Figure 14 shows an 8-cycle and 2-cycle variation on the 90-cycle periodicity. Its equation is

$$\overline{R}_{MAX} = 116.0 + 24.7 \sin \theta_5(X) + 20.3 \sin \theta_3(X) + 13.2 \sin \theta_4(X)$$
 (16)

Its 1-sigma spread is 30.4 and cycle 8 appears to be anomalously high.







Figure 13. 90 + 11 FIT.







Figure 14. 90 + 8 + 2 FIT.

Figure 15 depicts the function which had the smallest spread for the examples tested. It is an 11-cycle and 2-cycle variation on the 90-cycle periodicity. Its equation is:

$$\overline{R}_{MAX} = 116.0 + 24.7 \sin \theta_5(X) + 35.4 \sin \theta_2(X) + 15.9 \sin \theta_4(X)$$
 (17)

Its 1-sigma spread is 18.8 units, a 48-percent reduction from the baseline spread. (The variance has been reduced by 73 percent.) Cycle 17 appears to be anomalously low.







Figure 15. 90 + 11 + 2 FIT.

D. Pre-Cycle 8 Fits

According to the above analysis, values of \overline{R}_{MAX} for cycles 8 through 20, the most reliably known, can be fit very well by equation (17). Using this preferred empirical function, one can now extrapolate "backwards" in time to determine computed smoothed sunspot number at cycle maximum for earlier epochs. (It is realized that equation (17) is merely a fit for a small sample, and that this fit may have no "post-dictive" or "pre-dictive" value.) In Figure 16, \overline{R}_{MAX} is computed backwards in time to about A.D. 1080, corresponding to SCN -60. The top-most line is the 90 + 11 + 2 FIT and the bottom-most line is the 90 + 8 + 2 FIT, drawn for comparison. Observed values of \overline{R}_{MAX} for cycle 1 through 21 are drawn as open circles; computed values



computed $R_{\rm MAX}$ values for SCN -64 to SCN 30. Summary chart: Figure 16.

are drawn as darkened triangles. The approximate dates, as stated by Eddy [3,4], for the Maunder Minimum, Spörer Minimum, and Grand Maximum are shown. A table legend is included which identifies some major milestones in solar research, as related to the study of sunspots. Computed mean values and standard deviations are shown for the two minima and single maximum periods. A note reminding us that the best-quality sunspot data extends from about cycle 8 to the present is also included. One quickly observes that, while the 90 + 11 + 2 FIT gives a very good representation of cycles 8 through 21, it gives a very poor match to cycles 2 through 7, sometimes departing by more than 3 sigma from computed values. Cycles 2 through 7 appear to be better-fitted by the 90 + 8 + 2 FIT. This seems to be supportive of the belief that there is an 8-cycle periodicity imbedded in the sunspot record; however, how much weight should be given to this view and even to the firmness of the \overline{R}_{MAX} values for cycles 2 through 7 remain unclear. Additional investigation should be pursued.

Table 3 gives values for several selected cycle-related parameters for cycles 1 through 7. Mean values and standard deviations are given for the group cycles 1 through 7 and for the groups cycles 8 through 20 and 1 through 20. Clearly, when parametric values are compared for the different groupings, it is found that in some cases they are quite dissimilar; e.g., ASC and DES. This may be an indication that smoothed sunspot number at cycle maximum for cycles observed prior to cycle 8 are in error. (It is known from Waldmeier [170] that daily records of sunspot number, which form the basis for a determination of monthly mean sunspot number which is then used to compute smoothed sunspot number, are incomplete.) Rather, it may mean that the period encompassing cycles 2 through 7 was an anomalous timespan, with values departing from computed values by substantial amounts. (Very recently, Sonett [36] has addressed this particularly anomalous era and found the sunspot record to be consistent with the idea that, perhaps, there may have been a "book-keeping" error in the compilation of sunspot number about 1780 to 1800.)

TABLE 3. SELECTED PARAMETRIC VALUES FOR CYCLES 1 THROUGH 7

SCN	MINIMUM OCCURRENCE DATE	MAXIMUM OCCURRENCE DATE	Ř _{MIN}	R MAX	ASC	DES	MIN-MIN PERIOD
1	FEB 1755	JUN 1761	8.4	86.5	75	60	135
2	MAY 1766	SEP 1769	11.2	115.8	39	69	108
.3	JUN 1775	MAY 1778	7.2	158.5	35	76	111
4	SEP 1784	FEB 1788	9.5	141.2	41	96	137
5	APR 1798	FEB 1805	3.2	49.2	81	65	146
6	JUL 1810	MAY 1816	0.0	48.7	70	83	153
7	APR 1823	NOV 1829	0.1	71.7	79	48	127
		MEAN _{1—7} ^{\$} 1—7	5.7 4.2	95.9 40.4	60.0	71.0 14.6	131.0 15.6
		MEAN ₈₋₂₀	5.2	116.2	48.2	83.5	131.6
		^{\$} 8-20	2.7	36.7	6.2	10.0	10.0
		MEAN ₁₋₂₀	5.4	109.1	52.3	79.1	131.4
		^{\$} 1—20	3.3	39.2	13.6	13.2	12.2

Of the three aforementioned, preceding epochs, only the Maunder Minimum can be examined more closely. According to equation (17), as Figure 16 shows, a minimum $\overline{R}_{\rm MAX}$ value occurs in the 17th century. However, the values postdicted by equation (17) are higher than has been suggested for this period (e.g., Eddy [1,8]). (Equation (17) suggests a mean value for $\overline{R}_{\rm MAX}$ of about 91 \pm 32.) Recently, Xu and Jiang [35] re-assessed the 17th Century Chinese sunspot records and concluded that the inference to a Maunder Minimum may be incorrect (Link [23] and Schove [24]). They found sightings which allowed them to argue that solar cycles were continuing during the Maunder Minimum and that the average annual sunspot number at cycle maximum during this period was around 50 or less. Very recently, Zhang [37], based on Korean auroral records for the period A.D. 1507-1747, reports that solar activity during the Maunder Minimum may not have been as low as Eddy has claimed. Also, Hameed and Wyant [38] have found an approximate 23-year cycle in surface temperatures in central England during and after the Maunder Minimum, suggesting that if variation in climate may be partly due to solar activity, then solar activity may still have been very much apparent even during the Maunder Minimum. Figure 16 gives $\overline{R}_{\rm MAX}$ values which can be converted to annual numbers for direct comparison. Figure 17 plots annual sunspot number at cycle maximum, R(A), versus $\overline{R}_{\rm MAX}$




Figure 17. R(A) versus R_{MAX}.

based on cycles 8 through 20. The Pearson correlation coefficient r is 1.00 with a standard error of estimate S_{yx} equal to about 3.8. The equation relating the two parameters is:

$$R(A) = -1.063 + 0.952 \overline{R}_{MAX}$$
 (18)

Based on equation (18) and the 90 + 11 + 2 FIT, R(A) values are expected to have been about 86 ± 31 during the Maunder Minimum. If the Maunder Minimum is viewed as an anomalous timespan where R(A) values are depressed about 1 or more standard deviations below normal, then R(A) is expected to often be below 55, very near values reported by Xu and Jiang.

In Table 4, the Xu and Jiang results are recast along with those reported by Schove [22] and R(A) estimates based on the 90 + 11 + 2 and 50 + 11 + 2 FITS. The table spans SCN -14 to SCN -3, where the Maunder Minimum approximately spans SCN -9 to SCN -3. Six of the ten records (Xu and Jiang only give results for SCN -14 to SCN -5) for which Xu and Jiang give results are very compatible with those obtained using either the 90 + 11 + 2 or 50 + 11 + 2 FITS. If the statistical spread is allowed to be greater than 1-sigma, then, essentially all the records overlap. This comparison suggests that while a Maunder Minimum very well may have occurred, solar periodicities were probably still working and values of \overline{R}_{MAX} given by the empirical function (depressed by at least 1-sigma, possibly more) may be representative of the period. (Recently, Eddy [8] has re-apprecised the Maunder Minimum and

empirical function (depressed by at least 1-sigma, possibly more) may be representative of the period. (Recently, Eddy [8] has re-appraised the Maunder Minimum and argues that the basis for the occurrence of such a minimum remains firm, with peak annual sunspot number being of the order of 20 or below during this protracted period of low sunspot number.)

TABLE 4. COMPARISON OF REPORTED AND COMPUTED $\overline{R}_{\text{MAX}}$ VALUES FOR THE MAUNDER MINIMUM

SCN	R(A) ¹ est	R(A) ² est	R(A) ³ COMP	R(A) ⁴ COMP	NOTES
-14	50	80	106	102	
-13	>160	90	129	125	
-12	85	100	85	80	
-11	160	70	97	91	
-10	50	40	48	44	
- 9	50	(50	68	63	
8	45	{	38	33	1
– 7	< 45	60	79	71	
- 6	45	50	66	.58	MAUNDER MINIMUM
- 5	< 45	30	116	105	
- 4	*	50	100	89	
- 3	*	130	135	125	

^{*}No value given in Xu and Jiang [35]

¹According to Xu and Jiang [35], based on naked-eye Chinese sunspot records

According to Schove [22] Based on 90 + 11 + 2 FIT

⁴Based on 50 + 11 + 2 FIT

E. Cycles Beyond SCN 20

This report will now extrapolate the empirical function "forwards" in time, at least for cycles 21 and 22, the two cycles of more immediate concern. The 90 + 11 + 2 FIT predicts that cycle 21 has a 1-sigma confidence \overline{R}_{MAX} value equal to 157.3 \pm 18.8. This value is remarkably consistent with that predicted by Sargent [64] (= 154) and is very close to that which has now been observed (= 164.5). Table 5 lists estimates for \overline{R}_{MAX} for cycle 21 based on all the approaches given earlier; it also gives residuals. It was seen that cycle 21 was about 0.38 s unit above the predicted value, in the average range of Figure 15. Table 6 lists the cycle-related parameters projected for cycle 21, based on the 90 + 11 + 2 FIT and on the actually observed \overline{R}_{MAX} value. (Values for the cycle-related parameters are based on regression equations using \overline{R}_{MAX} as the independent variable, given in Figures 18 through 20, included in the Appendix.) Also given in Table 6 are the occurrence dates for \overline{R}_{MAX} , \overline{R}_{CHM} (ascent and descent side), and cycle end date, denoted here as \overline{R}_{MIN} (SCN + 1) occurrence which marks the beginning of the next cycle (SCN 22).

TABLE 5. \overline{R}_{MAX} (COMP) VALUES FOR SCN 21

METHOD	R _{MAX} (COMP) ± 1 SIGMA	RES = R _{MAX} (OBS) - R _{MAX} (COMP)
MEAN	116.2 ± 36.7	48.3
LR	130.1 ± 36.1	34.4
LR + 8	130.1 ± 33.4	34.4
LR + 11	142.8 ± 25.2	21.7
LR + 11 + 8	142.8 ± 24.9	21.7
LR + 11 + 2	158.6 ± 19.7	5.9
50	118.8 ± 37.0	45.7
50 + 8	118.8 ± 33.8	45.7
50 + 11	132.6 ± 24.5	31.9
50 + 11 + 8	132.6 ± 23.8	31.9
50 + 8 + 2	131.6 ± 31.2	32.9
50 + 11 + 2	148.0 ± 19.0	16.5
90	126.8 ± 36.1	37.7
90 + 8	126.8 ± 33.1	37.7
90 + 11	141.5 ± 24.6	23.0
90 + 8 + 2	140.0 ± 30.4	24.5
90 + 11 + 2	157.3 ± 18.8	7.2

Table 6 reveals that, based on the 90 + 11 + 2 FIT \overline{R}_{MAX} projection (= 157.3), cycle 21 maximum would have been estimated to have occurred about February 1980 and \overline{R}_{CHM} occurrence dates about October 1978 and January 1983; cycle end date would be expected to occur about March 1987. Based on actual dates of occurrence, it is noted that \overline{R}_{MAX} occurrence would have been missed by only 2 months (December 1979), and \overline{R}_{CHM} ascent-side and descent-side occurrences by 4 months and 2 months, respectively (June 1978 and March 1983). Cycle 21 is still in decline, so a comparison between computed and observed cycle end dates is not yet possible. Wilson [44] and Wilson et al. [45,100] have discussed the expected cycle duration for cycle 21. If cycle 21 is a short-period cycle (cycle duration less than 128 months),

TABLE 6. SELECTED PARAMETRIC VALUES FOR SCN 21

PARAMETER	MEAN VALUE ¹	STANDARD DEVIATION ¹	OBS VALUE	COMP. VALUE ²	COMP. VALUE ³
R _{MAX}	116.2	36.7	164.5	157.3	
R _{MIN}	5.2	2.7	12.2	6.1	6.2
R _{MEAN}	53.4	16.2		70.7	73.7
^S MEAN	35.9	12.2		48.3	50.4
^R CHM	60.7	18.8	88.4	81.7	85.4
ASC _{CHM}	26.3	5.8	24	27.6	27.8
D _{CHM}	56.1	9.2		50.9	50.0
ASC	48.2	6.2	42	44.3	43.6
DES	83.5	10.0		84.5	84.7
MIN-MIN PÉRIOD	131.6	10.0		128.8	128.3
SLOPEASC	2.398	0,958	3.626	3.426	3.606
SLOPEDES	-1.337	0.441		-1.767	-1.851
, , , , , , , , , , , , , , , , , , ,	water and the control of the control		OBS.	COMP.2	COMP.3
R _{MAX} OCCURRE	NCE DATE:		DEC 1979	FEB 1980	
R _{CHM} OCCURRE	NCE DATES:	ASC-SIDE	JUN 1978	OCT 1978	OCT 1978
R _{MIN} (SCN + 1) O	CCURRENCE DATE	DES-SIDE	MAR 1983	JAN 1983 MAR 1987	DEC 1982 FEB 1987

¹BASED ON CYCLES 8 THROUGH 20

then cycle 21 should end before March 1987 (best-guess date: August 1986); otherwise, cycle 21 will be a long-period cycle (cycle duration greater than 133 months) ending no earlier than August 1987 (best-guess date: February 1988).

In Table 7, computed \overline{R}_{MAX} values with 1-sigma confidence for cycle 22 are listed. Extrapolation of the 90 + 11 + 2 FIT yields an estimate of \overline{R}_{MAX} for cycle 22 to be 107.4. (Sargent and Heckman [66] and Sargent [171] have discussed current efforts to predict cycle 22; also, a Solar Terrestrial Prediction Workshop with the expressed objective of discussing current efforts to predict cycle 22 occurred 18-22 June 1984 in Meudon, France.) Using this value, Table 8 projects values for the cycle-related parameters for cycle 22. The \overline{R}_{MAX} occurrence date is projected to be

 $^{^{2}}$ BASED ON 90 + 11 + 2 FIT; \overline{R}_{MAX} (COMP) = 157.3

 $^{^{3}}$ BASED ON \tilde{R}_{MAX} (OBS) = 164.5

TABLE 7. $\overline{R}_{\mbox{\scriptsize MAX}}$ (COMP) VALUES FOR SCN 22

METHOD	R _{MAX} (COMP) ± 1 SIGMA	
LR + 11 + 2 FIT	111.9 ± 19.7	
50 + 11 + 2 FIT	98.0 ± 19.0	
90 + 11 + 2 FIT	107.4 ± 18.8	
90 + 8 + 2 FIT	100.8 ± 24.5	

TABLE 8. SELECTED PARAMETRIC VALUES FOR SCN 22

PARAMETER	MEAN VALUE ¹	STANDARD DEVIATION ¹	COMPUTED VALUE
\bar{R}_{MAX}	116.2	36.7	107.4
R _{MIN}	5.2	2,7	5.0
R MEAN	53.4	16.2	49.7
SMEAN	35.9	12.2	33.3
Ř _{CHM}	60.7	18.8	56.2
ASC _{CHM}	26.3	5.8	26.0
D _{CHM}	56.1	9.2	57.2
ASC	48.2	6.2	49.0
DES	83.5	10.0	83.2
MIN-MIN PERIOD	131.6	10.0	132.2
SLOPEASC	2.398	0.958	2.178
SLOPEDES	-1.337	0.441	-1.245
R _{MIN} OCCURRENCE DATE:		<u>COMP.</u> MAR 1987	
R _{MAX} OCCURRENCE DATE:		APR 1991	
R _{CHM} OCCURRENCE DATES: ASC-SIDE DES-SIDE		MAY 1989 FEB 1994	
R _{MIN} (SCN + 1) OCCURRENCE DATE:		MAR 1998	

¹BASED ON CYCLES 8 THROUGH 20

 $^{^2}$ BASED ON 90 + 11 + 2 FIT; \overline{R}_{MAX} (COMP) = 107.4

April 1991, \overline{R}_{CHM} occurrence dates May 1989 and February 1994, and the cycle-end date March 1998, all dates based on the March 1987 projected end date for cycle 21. Other dates, based on bifurcation of sunspot cycles into short-period and long-period cycles, have been summarized elsewhere (Wilson et al. [45,100]). In contrast to cycle 21 which is the second highest documented \overline{R}_{MAX} cycle, cycle 22 is projected to be a low-valued \overline{R}_{MAX} cycle, very probably of long cycle duration. Future cycles are projected to be decreasing in \overline{R}_{MAX} value through at least cycle 24 and then to be increasing in \overline{R}_{MAX} value thereafter, at least through cycle 30 (Fig. 16). The next "Grand Maximum" like period is projected to be around SCN 40, corresponding to ca. A.D. 2180.

IV. CONCLUSIONS

This study has shown that values of \overline{R}_{MAX} for modern sunspot cycles (i.e., cycles 8 through 21) can be satisfactorily fitted using a three-component sine curve representation, having an 11-cycle and a 2-cycle variation imbedded in an approximately 90-cycle periodicity (in terms of sunspot cycle number). The 90 + 11 + 2 FIT yields "minima," non-zero, low-value \overline{R}_{MAX} periods, such as the Maunder and Spörer Minima, and a "maximum," high-value $\overline{\overline{R}}_{ ext{MAX}}$ period, such as the Grand Maximum, as reported by Eddy [3,4], although the values of $\overline{R}_{ ext{MAX}}$ projected for the protracted sunspot minimum periods are considerably higher than has been estimated by other investigators [8,35]. Annual sunspot numbers, as reported by Xu and Jiang [35], for the Maunder Minimum are found to be very close to values "post-dicted" by the 90 + 11 + 2 FIT, especially if we consider the Maunder Minimum (and other such minima) to be a brief period when sunspot number is about 1-sigma below "normal" (i.e., below computed values). The 90 + 11 + 2 FIT, extrapolated forward in time, has been used to give estimates of \overline{R}_{MAX} for cycles 21 and 22, and a number of useful linear relationships based on $\overline{R}_{\mathrm{MAX}}$ [44] has been applied to project the behavior of these cycles. A comparison of observed and predicted values for the limited number of known cycle-related parameters for cycle 21 yields very good agreement. Cycle 22 is predicted to be a low-valued \overline{R}_{MAX} /long-period cycle, with \overline{R}_{MAX} equal to about 107.

APPENDIX — LINEAR REGRESSIONS BASED ON $\overline{R}_{ ext{MAX}}$




Figure 18. SLOPE ASC, \bar{R}_{CHM} , \bar{R}_{MEAN} and \bar{R}_{MIN} versus \bar{R}_{MAX} .




Figure 19. MIN-MIN PERIOD, D_{CHM} and ASC_{CHM} versus \overline{R}_{MAX} .




Figure 20. DES and ASC versus $\overline{\boldsymbol{R}}_{MAX}$




Figure 21. $SLOPE_{DES}$ versus $SLOPE_{ASC}$.




Figure 22. s_{MEAN} versus \overline{R}_{MEAN} .

REFERENCES

- 1. Eddy, J. A.: The Maunder Minimum. Science, Vol. 192, 1976, pp. 1189-1202.
- 2. Eddy, J. A.: The Sun Since the Bronze Age. Physics of Solar Planetary Environments, Proceedings of the International Symposium on Solar Terrestrial Physics, June 7-18, 1976, Boulder, Colorado, (ed., D. J. Williams), Vol. II, AGU, Boulder, Colorado, 1976, pp. 958-972.
- 3. Eddy, J. A.: Climate and the Changing Sun. Climatic Change, Vol. 1, 1977, pp. 173-190.
- 4. Eddy, J. A.: Historical Evidence for the Existence of the Solar Cycle. The Solar Output and Its Variation, (ed., O. R. White), Colorado Associated University Press, Boulder, Colorado, 1977, pp. 51-71.
- 5. Eddy, J. A.: The Case of the Missing Sunspots. <u>Sci. Am.</u>, Vol. 236, 1977, pp. 80-88, 92.
- 6. Eddy, J. A.: Historical and Aboreal Evidence for a Changing Sun. <u>The New Sun</u> (ed., J. A. Eddy), AAAS Selected Symp. 17, Westview Press, Boulder, Colorado, 1978, pp. 11-33.
- 7. Eddy, J. A.: The Historical Record of Solar Activity. The Ancient Sun: Fossil Record in the Earth, Moon and Meteorites (eds., R. O. Pepin, J. A. Eddy, and R. B. Merrill), Geochimica et Cosmochimica Acta Supplement 13, Pergamon Press, New York, New York, 1980, pp. 119-134.
- 8. Eddy, J. A.: The Maunder Minimum: A Reappraisal. Solar Phys., Vol. 89, 1983, pp. 195-207.
- 9. Stuiver, M. and Grootes, P. M.: Trees and the Ancient Record of Heliomagnetic Cosmic Ray Flux Modulation. The Ancient Sun: Fossil Record in the Earth, Moon and Meteorites (eds., R. O. Pepin, J. A. Eddy, and R. B. Merrill), Geochimica et Cosmochimica Acta Supplement 13, Pergamon Press, New York, New York, 1980, pp. 165-173.
- 10. Stuiver, M. and Quay, P. D.: Changes in Atmospheric Carbon-14 Attributed to a Variable Sun. Science, Vol. 207, 1980, pp. 11-19.
- 11. Stuiver, M.: Solar Variability and Climatic Change During the Current Millenium. Nature, Vol. 286, 1980, pp. 868-871.
- 12. Siscoe, G. L.: Evidence in the Auroral Record for Secular Solar Variability. Rev. Geophys. Space Sci., Vol. 18, 1980, pp. 647-658.
- 13. Schroder, W.: Auroral Frequency in the 17th and 18th Centuries and the "Maunder Minimum." J. Atmos. Terr. Phys., Vol. 41, 1979, pp. 445-446.
- 14. Link, F.: L'Activite Solaire au 17^{ème} Siècle. L'Astronomie, Vol. 91, 1977, pp. 191-199.
- 15. Cullen, C.: Was There a Maunder Minimum? Nature, Vol. 283, 1980, pp. 427-428.

- 16. Robinson, L. J.: No Maunder Minimum? Sky Telesc., Vol. 64, September 1982, p. 234.
- 17. Maunder, W.: A Prolonged Sun-Spot Minimum. Knowledge, Vol. 17, 1894, pp. 173-176.
- 18. Clerke, A.: A Prolonged Sun-Spot Minimum. Knowledge, Vol. 17, 1894, p. 206.
- 19. Clark, D. H. and Stephenson, F. R.: An Interpretation of the Pre-Telescopic Sunspot Periods from the Orient. Quart. J. Roy. Astron. Soc., Vol. 19, 1978, pp. 387-410.
- 20. Strothers, R.: Ancient Aurorae. Isis, Vol. 70, 1979, pp. 85-95.
- 21. Strothers, R.: Solar Activity Cycle During Classical Antiquity. <u>Astron.</u> Astrophys., Vol. 77, 1979, pp. 121-127.
- 22. Schove, D. J.: The Sunspot Cycle, 649 B.C. to A.D. 2000. <u>J. Geophys.</u> Res., Vol. 60, 1955, pp. 127-146.
- 23. Link, F.: Solar Cycles Between 1540 and 1700. Solar Phys., Vol. 59, 1978, pp. 175-178.
- 24. Schove, D. J.: Sunspot Turning-Points and Aurorae Since A.D. 1510. Solar Phys., Vol. 63, 1979, pp. 423-432.
- 25. Mitchell, J. M. Jr.: Evidence of a 22-Year Rhythm of Drought in the Western United States Related to the Hale Solar Cycle Since the 17th Century. Solar-Terrestrial Influence on Weather and Climate (eds., B. M. McCormac and T. A. Seliga), D. Reidel Publ. Co., Dordrecht, Holland, 1979, pp. 125-143.
- 26. Link, F.: Manifestations de l'Activite Solaire dans le Passe Historique. <u>Planet</u>. Space Sci., Vol. 12, 1964, pp. 333-348.
- 27. Link, F.: Solar Activity During the 17th Century. Astron. Astrophys., Vol. 54, 1977, pp. 857-861.
- 28. Wittman, A.: The Sunspot Cycle Before the Maunder Minimum. Astron. Astrophys., Vol. 66, 1978, pp. 93-97.
- 29. Vitinsky, Yu. I.: Comments on the So-Called Maunder Minimum. Solar Phys., Vol. 57, 1978, pp. 475-478.
- 30. Suess, S. T.: The Solar Wind During the Maunder Minimum. Planet. Space Sci., Vol. 27, 1979, pp. 1001-1013.
- 31. Schove, D. J.: Sunspot Turning-Points and Aurorae Since A.D. 1510. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A111-A128.
- 32. Robock, A.: Internally and Externally Caused Climate Change. <u>J. Atmos. Sci.</u>, Vol. 35, 1978, pp. 1111-1122.

- 33. Mass, C. and Schneider, S. H.: Statistical Evidence on the Influence of Sunspots and Volcanic Dust on Long-Term Temperature Records. <u>J. Atmos.</u> Sci., Vol. 34, 1977, pp. 1995-2004.
- 34. Morth, H. T. and Schlamminger, L.: Planetary Motion, Sunspots, and Climate. Solar-Terrestrial Influences on Weather and Climate (eds., B. M. McCormac and T. A. Seliga), D. Reidel Publ. Co., Dordrecht, Holland, 1979, pp. 193-207.
- 35. Xu Zhen-tao and Jiang Yao-tiao: The Solar Activity in the Seventeenth Century Re-Assessed in the Light of Sunspot Records in the Local Gazettes of China. Chin. Astron. Astrophys., Vol. 6, 1982, pp. 84-90.
- 36. Sonett, C. P.: The Great Solar Anomaly ca. 1780-1800: An Error in Compiling the Record? J. Geophys. Res., Vol. 88, 1983, pp. 3225-3228.
- 37. Zhang, Z.: The Korean Auroral Records of the Period A.D. 1507-1747 and the SAR Arcs. SUIPR Report No. 955, Institute for Plasma Research, Stanford, California, April 1983 (submitted to J. Geophys. Res.).
- 38. Hameed, S. and Wyant, P.: Twenty-Three Year Cycle in Surface Temperatures During the Maunder Minimum. Geophys. Res. Letters, Vol. 9, 1982, pp. 83-86.
- 39. Ding You-ji, Luo Bao-rong and Feng Yong-ming: Peak Years of Various Solar Cycles. Chin. Astron. Astrophys., Vol. 7, 1983, pp. 24-30.
- 40. Clark, D. and Stephenson, F. R.: Establishing the Case for the Maunder Minimum. Nature, Vol. 284, 1980, p. 592.
- 41. Willis, D. M., Easterbrook, M. G. and Stephenson, F. R.: Seasonal Variation of Oriental Sunspot Sightings. Nature, Vol. 287, 1980, pp. 617-619.
- 42. Landsberg, H. E.: Arch. Met. Geoph. Biokl. Ser. B., Vol. 28, 1980, p. 181.
- 43. Link, F.: Auroral and Climatic Cycles in the Past. J. Brit. Astron. Assoc., Vol. 78, 1968, pp. 195-205.
- 44. Wilson, R. M.: A Comparative Look at Sunspot Cycles. NASA TP-2325, Marshall Space Flight Center, Alabama, May 1984.
- 45. Wilson, R. M., Reichmann, E. J. and Teuber, D. L.: An Empirical Method for Estimating Sunspot Number. Contribution to the Solar-Terrestrial Prediction Workshop (18-22 June 1984; Meudon, France), paper No. A.1.8 Long Term Solar Prediction, in press.
- 46. Wallenhorst, S. G.: Sunspot Numbers and Solar Cycles. Sky <u>Telesc.</u>, Vol. 62, September 1982, pp. 234-236.
- 47. Newkirk, G. Jr. and Frazier, K.: The Solar Cycle. Phys. Today, Vol. 35, April 1982, pp. 25-34.
- 48. Bray, R. J. and Loughhead, R. E.: Sunspots. John Wiley and Sons, Inc., New York, New York, 1964.

- 49. Noyes, R. W.: <u>The Sun</u>, <u>Our Star</u>. Harvard University Press, Cambridge, Massachusetts, 1982.
- 50. Meadows, A. J.: Early Solar Physics. Pergamon Press, London, England, 1970.
- 51. Rubashev, B. M.: Problems of Solar Activity. NASA TT F-244, Washington, D.C., 1964.
- 52. Vitinskii, Yu. I.: Solar-Activity Forecasting. NASA TT F-289, Washington, D.C., 1965.
- 53. Bonov, A. D.: Characteristic Features of Secular and Supersecular Cycles of Solar Activity. A 180-Year Cycle. Solar Activity and Related Interplanetary and Terrestrial Phenomena (ed., J. Xanthakis), Springer-Verlag, Berlin, Germany, 1973, pp. 83-86.
- 54. Bumba, V. and Kleczek, J. (eds.): <u>Basic Mechanisms of Solar Activity</u>. IAU Symp. 71, D. Reidel Publ. Co., Dordrecht, Holland, 1976.
- 55. Herman, J. R. and Golberg, R. A.: Sun, Weather and Climate. NASA SP-426, Washington, D.C., 1978, pp. 12-24.
- 56. Lundquist, C. A. and Vaughan, W. W.: Analysis of Regression Methods for Solar Activity Forecasting. Solar-Terrestrial Influences on Weather and Climate (eds., B. M. McCormac and T. A. Seliga), D. Reidel Publ. Co., Dordrecht, Holland, 1979, pp. 59-64.
- 57. Brier, G. W.: Use of Difference Equation Methods for Predicting Sunspot Numbers. Solar-Terrestrial Influences on Weather and Climate (eds., B. M. McCormac and T. A. Seliga), D. Reidel Publ. Co., Dordrecht, Holland, 1979, pp. 209-214.
- 58. Barnes, J.A., Sargent, H. H. III and Tryon, P. V.: Sunspot Cycle Simulation Using Random Noise. The Ancient Sun: Fossil Record in the Earth, Moon and Meteorites (eds., R. O. Pepin, J. A. Eddy, and R. B. Merrill), Geochimica et Cosmochimica Acta Supplement 13, Pergamon Press, New York, New York, 1980, pp. 159-163.
- 59. Romanchuk, P. R.: The Nature of Sunspot Cyclicity. I. <u>Sov. Astron.</u>, Vol. 25, 1981, pp. 87-92.
- 60. de Meyer, F.: Mathematical Modelling of the Sunspot Cycle. Solar Phys., Vol. 70, 1981, pp. 259-272.
- 61. Holland, R. L., Rhodes, C. A. and Euler, H. C. Jr.: Lagrangian Least-Squares Prediction of Solar Activity. NASA TM-82462, Marshall Space Flight Center, Alabama, April 1982.
- 62. Wallenhorst, S. G.: On Long-Term Periodicities in the Sunspot Cycle. <u>Solar</u> Phys., Vol. 80, 1982, pp. 379-384.
- 63. Zhukov, L. V. and Muzalevskii, Yu. S.: A Correlation Spectral Analysis of the Periodicities in Solar Activity. Sov. Astron., Vol. 13, 1969, pp. 473-479.

- 64. Sargent, H. H.: A Prediction for the Next Sunspot Cycle. <u>IEEE Vehicular Technology Conference</u>, 1978, pp. 490-496.
- 65. Simon, P. A.: Polar Coronal Holes and Solar Cycles. Solar Phys., Vol. 63, 1979, pp. 399-410.
- 66. Sargent, H. H. III and Heckman, G. R.: An Early Forecast of the Size of Suspot Cycle 22. <u>EOS</u>, Vol. 63, 1982, p. 1089.
- 67. Dicke, R. H.: Is There a Chronometer Hidden Deep in the Sun? Nature, Vol. 276, 1978, pp. 676-680.
- 68. Dicke, R. H.: Solar Luminosity and the Sunspot Cycle. Nature, Vol. 280, 1979, pp. 24-27.
- 69. Dale, J. B.: The Fluctuations in the Epochs of Sun-Spot Maxima and Minima. Mon. Not. Roy. Astron. Soc., Vol. 74, 1914, pp. 664-669.
- 70. Stewart, J. Q. and Panofsky, H. A. A.: The Mathematical Characteristics of Sunspot Variations. Astrophys. J., Vol. 88, 1938, pp. 385-407.
- 71. Stewart, J. Q. and Eggleston, F. C.: The Mathematical Characteristics of Sunspot Variations. II. Astrophys. J., Vol. 91, 1940, pp. 72-82.
- 72. Gleissberg, W.: Probability Laws of Sunspot Variations. Astrophys. J., Vol. 96, 1942, pp. 234-238.
- 73. Gleissberg, W.: The Predictability of the Probable Features of the Sunspot Cycles. Astrophys. J., Vol. 109, 1949, pp. 321-326. (Erratum, Astrophys. J., Vol. 109, 1949, p. 555.)
- 74. Gleissberg, W.: The Probable Behavior of the Next Sunspot Cycle. Astrophys. J., Vol. 110, 1949, pp. 90-95.
- 75. Cook, A. F.: On the Mathematical Characteristics of Sunspot-Variations. J. Geophys. Res., Vol. 54, 1949, pp. 347-354.
- 76. Granger, C. W. J.: A Statistical Model for Sunspot Activity. Astrophys. J., Vol. 126, 1957, pp. 152-158.
- 77. Ol', A. I.: The Possible Periodicity of Solar Activity. Sov. Astron., Vol. 4, 1960, pp. 212-216.
- 78. Godoli, G. and Allen, C. W.: The Indices of Solar Activity. Planet. Space Sci., Vol. 12, 1964, pp. 349-354.
- 79. Gleissberg, W.: Secularly Smoothed Data on the Minima and Maxima of Sunspot Frequency. Solar Phys., Vol. 2, 1967, pp. 231-233.
- 80. Zhukov, L. V. and Muzalevskii, Yu. S.: Comparison of the Observed Monthly Wolf Numbers as Smoothed by the Moving-Average and Whittaker Operators. Sov. Astron., Vol. 14, 1970, pp. 287-300.
- 81. Muzalevskii, Yu. S. and Zhukov, L. V.: The Fluctuations in the Monthly Wolf Numbers. Sov. Astron., Vol. 14, 1970, pp. 432-439.

- 82. Henkel, R.: A Feature of the Secularly Smoothed Maxima of Sunspot Frequency. Solar Phys., Vol. 20, 1971, pp. 345-347.
- 83. Hartmann, R.: A New Representation of the 80-Year Cycle in Sunspot Frequency. Solar Phys., Vol. 21, 1971, pp. 246-248.
- 84. Cole, T. W.: Periodicities in Solar Activity. Solar Phys., Vol. 30, 1973, pp. 103-110.
- 85. Gleissberg, W.: Revision of the Probability Laws of Sunspot Variations. Solar Phys., Vol. 30, 1973, pp. 539-542.
- 86. Bhargava, B. N. and Rangarajan, G. K.: Recurrent Magnetic Activity, Sunspot Number and Its Rate of Decline. Solar Phys., Vol. 40, 1975, pp. 235-239.
- 87. Wolff, C. L.: Timing of Solar Cycles by Rigid Internal Rotations. Astrophys. J., Vol. 205, 1976, pp. 612-621.
- 88. Gnevyshev, M. N.: Essential Features of the 11-Year Solar Cycle. Solar Phys., Vol. 51, 1977, pp. 175-183.
- 89. Xanthakis, J. and Poulakos, C.: A Forecast of Solar Activity for the 21st Solar Cycle. Solar Phys., Vol. 56, 1978, pp. 467-469.
- 90. Yoshimura, H.: The Solar-Cycle Period-Amplitude Relation as Evidence of Hysteresis of the Solar-Cycle Nonlinear Magnetic Oscillation and the Long-Term (55 Year) Cyclic Modulation. Astrophys. J., Vol. 227, 1979, pp. 1047-1058.
- 91. Piddington, J. H.: Theory of the Solar 22-Year Cycle. Proc. ASA, Vol. 2, 1971, pp. 7-10.
- 92. Bray, J. R.: A Sunspot-Auroral Solar Index from 522 B.C. to A.D. 1968. New Zealand J. Sci., Vol. 23, 1980, pp. 99-106.
- 93. Kane, R. P. and Trivedi, N. B.: Relationship Between Sunspot Numbers During Years of Sunspot Maximum and Sunspot Minimum. Solar Phys., Vol. 68, 1980, pp. 135-139.
- 94. Yallop, B. D. and Hohenkerk, C. Y.: Distribution of Sunspots 1874-1976. Solar Phys., Vol. 68, 1980, pp. 303-305.
- 95. Brown, G. M. and Butcher, E. C.: The Use of Abnormal Quiet Days in Sq(H) for Predicting the Magnitude of Sunspot Maximum at the Time of Preceding Sunspot Minimum. Planet. Space Sci., Vol. 29, 1981, pp. 73-77.
- 96. Legrand, J. P. and Simon, P. A.: Ten Cycles of Solar and Geomagnetic Activity. Solar Phys., Vol. 70, 1981, pp. 173-195.
- 97. Waldmeier, M.: The Parameters of Solar Cycle No. 21. Solar Phys., Vol. 73, 1981, p. 207.
- 98. Voichishin, K. S. and Stodilka, M. I.: On the Statistical Stability of Solar Cyclicity. Sov. Astron., Vol. 26, 1982, pp. 705-711.

- 99. Kapoor, S. G. and Wu, S. M.: Developing Forecasting Charts for Sunspot Numbers. J. Geophys. Res., Vol. 87, 1982, pp. 9-16.
- 100. Wilson, R. M., Rabin, D. and Moore, R. L.: Bi-Modality of the Solar Cycle. Science, 1984 (in preparation).
- 101. Holland, R. L. and Vaughan, W. W.: Lagrangian Least-Squares Prediction of Solar Flux ($\overline{F}_{10.7}$). J. Geophys. Res., Vol. 89, 1984, pp. 11-16.
- 102. Wilson, R. M.: Sunspot Variation and Selected Associated Phenomena: A Look at Solar Cycle 21 and Beyond. NASA TM-82474, Marshall Space Flight Center, Alabama, February 1982.
- 103. Jose, P. D.: Sun's Motion and Sunspots. <u>Astron. J.</u>, Vol. 70, 1965, pp. 193-200.
- 104. Fairbridge, R. W. and Hameed, S.: Phase Coherence of Solar Cycle Minima over Two 178-Year Periods. Astron. J., Vol. 88, 1983, pp. 867-869.
- 105. Otaola, J. A. and Zenteno, G.: On the Existence of Long-Term Periodicities in Solar Activity. Solar Phys., Vol. 89, 1983, pp. 209-213.
- 106. Gregg, D. P.: A Nonlinear Solar Cycle Model with Potential for Forecasting on a Decadel Time Scale. Solar Phys., Vol. 90, 1984, pp. 185-194.
- 107. Teuber, D. L., Reichman, E. J. and Wilson, R. M.: Description of Sunspot Cycles by Orthogonal Functions. Astron. Astrophys., 1984, in press.
- 108. Solar Activity Prediction Group (Peking Observatory): Solar Activity Predictions at Peking Observatory. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 1, Boulder, Colorado, August 1979, pp. 154-162.
- 109. Xu Zhen-tao, Zhao Ai-di, Mei Yan-lin and Guo Quan-shi: Long-Term Forecasting of Solar Activity. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 1, Boulder, Colorado, August 1979, pp. 163-167.
- 110. Fougere, P. F.: Sunspots: Power Spectra and a Forecast. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A1-A9.
- 111. Vasilyev, O. B. and Kandaurova, K. A.: A Statistical Model of Solar Periodicity and Its Prediction. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A10-A21.
- 112. Hunter, H. E.: Application of an Eigenvector Technique to Prediction of Solar Activity. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A22-A36.
- 113. Wood, C. A.: Solar Activity Prediction for Cycle 21. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A37-A40.
- 114. Schatten, K. H.: Babcock Dynamo Theory and the Maunder Minimum. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A41-A44.

- 115. Vassilyea, G. I., Schpitalnaya, A. A. and Petrova, N. S.: On the Prediction of Solar Activity Taking into Account Its Extraneous Conditionality. Solar—Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A45-A57.
- 116. Feldstein, Y. I., Livshits, M. A. and Valtchuk, T. E.: Forecasting of Solar and Geomagnetic Activity Based on the Relation Between Geomagnetic Disturbance and High-Latitude Magnetic Field of the Sun. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A58-A70.
- 117. Vitinsky, Yu. I. and Rubashev, B. M.: Modification of Methods for Forecasting Monthly and Quarterly Wolf Numbers. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A71-A76.
- 118. Xanthakis, J. N. and Poulakos, C.: Prediction of the Radio Emission Indices of the Sun in the Frequency Range 1000 ≤ F ≤ 3750 MHz. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A77-A95.
- 119. Suda, T.: An Attempt at Prediction of Solar Activity Indices Which are Closely Related to Climate. Solar-Terrestrial Predictions Proceedings (ed., R. F. Donnelly), Vol. 3, Boulder, Colorado, March 1980, pp. A96-A110.
- 120. Cohen, T. J. and Lintz, P. R.: Long Term Periodicities in the Sunspot Cycle.

 Nature, Vol. 250, 1974, pp. 398-400.
- 121. Wood, K. D.: Sunspots and Planets. Nature, Vol. 240, 1972, pp. 91-93.
- 122. Antalova, A. and Gnevyshev, M. N.: Principal Characteristics of the 11-Year Solar Activity Cycle. <u>Sov. Astron.</u>, Vol. 9, 1965, pp. 198-201.
- 123. McNish, A. G. and Lincoln, J. V.: Prediction of Sunspot Numbers. <u>Trans.</u> Am. Geophys. Un., Vol. 30, 1949, pp. 673-685.
- 124. Suda, T.: Some Statistical Aspects of Solar-Activity Indices. <u>J. Met. Soc.</u> <u>Japan</u>, Vol. 40, 1962, pp. 287-299.
- 125. King-Hele, D. G.: Prediction of the Dates and Intensities of the Next Two Sunspot Maxima. Nature, Vol. 209, 1966, pp. 285-286.
- 126. Sleeper, H. P. Jr.: Solar Activity Prediction Methods. Northrup Report TR-230-1405, Huntsville, Alabama, January 1975 (NAS8-21810).
- 127. Sleeper, H. P. Jr.: Planetary Resonances, Bi-Stable Oscillation Modes and Solar Activity Cycles. Northrup Report TR-241-1053, Huntsville, Alabama, February 1972 (NASA CR-2035).
- 128. Sleeper, H. P. Jr.: Bi-Stable Oscillation Modes of the Sun and Long Range Prediction of Solar Activity. Northrup Report TR-709, Huntsville, Alabama, February 1970 (NAS8-20082).
- 129. Euler, H. C. Jr.: High Resolution Power Spectra of Daily Zurich Sunspot Numbers. NASA TM X-64718, Marshall Space Flight Center, Alabama, 1973.

- 130. Yule, G. U.: VII. On a Method of Investigating Periodicities in Distributed Series, with Special Reference to Wolfer's Sunspot Numbers. Phil. Trans. Roy. Soc. (Ser. A), Vol. 226, 1926-27, pp. 267-298.
- 131. Stewart, J. Q. and Eggleston, F. C.: A Prediction of the Remaining Course of the Present Sunspot Cycle. Phys. Rev., Vol. 55, 1939, p. 1102.
- 132. Schuster, A.: On the Periodicity of Sun-Spots. Proc. Roy. Soc. London (Ser. A), Vol. 85, 1911, pp. 50-53.
- 133. Schuster, A.: The Influence of Planets on the Formation of Sun-Spots. Proc. Roy. Soc. London (Ser. A), Vol. 85, 1911, pp. 309-323.
- 134. Bell, B. and Wolbach, L. G.: Lunar Eclipses and the Forecasting of Solar Minima. Icarus, Vol. 4, 1965, pp. 409-414.
- 135. Bigg, E. K.: Influence of the Planet Mercury on Sunspots. Astron. J., Vol. 72, 1967, pp. 463-466.
- 136. Wood, R. M. and Wood, K. D.: Solar Motion and Sunspot Comparison. Nature, Vol. 208, 1965, pp. 129-131.
- 137. King-Hele, D. G.: Prediction of Future Sunspot Cycles. <u>Nature</u>, Vol. 199, 1963, pp. 226-227.
- 138. Xanthakis, J.: Expression de l'Activite Solaire en Fonction du Temps d'Ascension. Advances in <u>Upper Atmosphere Research</u> (ed., B. Landmark), MacMillan Co., New York, New York, 1963, pp. 285-302.
- 139. Westcott, P.: The 25- or 26-Month Periodic Tendency in Sunspots. <u>J. Atmos.</u> Sci., Vol. 21, 1964, pp. 572-573.
- 140. Shapiro, R. and Ward, F.: A Neglected Cycle in Sunspot Numbers? J. Atmos. Sci., Vol. 19, 1962, pp. 506-508.
- 141. Minnis, C. M.: An Estimate of the Peak Sunspot Number in 1968. <u>J. Atmos.</u> Terr. Phys., Vol. 20, 1961, pp. 94-99.
- 142. Huttly, N. A.: A Review of Recent Investigations into Sun-Spot Cycles. Marconi Rev., Vol. 20, 1957, pp. 117-129.
- 143. Anderson, C. N.: Notes on the Sunspot Cycle. <u>J. Geophys. Res.</u>, Vol. 59, 1954, pp. 455-461.
- 144. Gleissberg, W.: The Eighty-Year Solar Cycle in Auroral Frequency Numbers.

 J. Brit. Astron. Ass., Vol. 75, 1965, pp. 227-231.
- 145. Shapley. A. H.: An Estimate of the Trend of Solar Activity, 1944-1950. J. Terr. Mag. Elec., Vol. 49, 1944, pp. 43-45.
- 146. Clayton, H.: The Sunspot Period. Smithsonian Misc. Coll., Vol. 98, No. 2, 1939.
- 147. Linder, R. C.: Prediction of Next Sunspot Maximum. Pop. Astron., Vol. 53, 1945, p. 250.

- 148. Maunder, E. W.: The Prolonged Sunspot Minimum, 1645-1715. <u>J. Brit. Astron.</u>
 <u>Ass.</u>, Vol. 32, 1922, pp. 140-145.
- 149. Anderson, C. N.: A Representation of the Sunspot Cycle. Terr. Mag., Vol. 44, 1939, pp. 175-179.
- 150. Schove, D. J.: Sunspot Cycles. Hutchinson Ross, Stroudsburg, Benchmark, Vol. 68, 1983.
- 151. Gleissberg, W.: The Eighty-Year Sunspot Cycle. J. Brit. Astron. Ass., Vol. 68, 1958, pp. 148-152.
- 152. Herrinck, P.: Prediction of Sunspot Numbers Until the End of the Present Cycle. Nature, Vol. 184, 1959, pp. 51-52.
- 153. Okal, E. and Anderson, D. L.: On the Planetary Theory of Sunspots. Nature, Vol. 253, 1975, pp. 511-513.
- 154. Kane, R. P.: Predicted Intensity of the Solar Maximum. Nature, Vol. 274, 1978, pp. 139-140.
- 155. Bracewell, R. N.: The Sunspot Number Series. Nature, Vol. 171, 1953, pp. 649-650.
- 156. Currie, R. G.: Fine Structure in the Sunspot Spectrum 2 to 70 Years. Astrophys. Space Sci., Vol. 20, 1973, pp. 509-518.
- 157. Radowski, H. R., Fougere, P. F. and Zawalick, E. J.: A Comparison of Power Spectral Estimates and Applications of the Maximum Entropy Method. J. Geophys. Res., Vol. 80, 1975, pp. 619-625.
- 158. Williams, D.: Sunspot Cycle Correlations. Ann. N. Y. Acad. Sci., Vol. 95, 1961, pp. 78-88.
- 159. Schove, D. J.: Solar Cycles and the Spectrum of Time Since 200 B.C. Ann. N. Y. Acad. Sci., Vol. 95, 1961, pp. 107-123.
- 160. Brown, G. M.: What Determines Sunspot Maximum? Mon. Not. Roy. Astron. Soc., Vol. 174, 1976, pp. 185-189.
- 161. Schatten, K. H., Scherrer, P. H., Svalgaard, L. and Wilcox, J. M.: Using Dynamo Theory to Predict the Sunspot Number During Solar Cycle 21. Geophys. Res. Letters, Vol. 5, 1978, pp. 411-414.
- 162. Kondratyev, K. Ya. and Nikolsky, G. A.: Solar Radiation and Solar Activity. Quart. J. Roy. Met. Soc., Vol. 96, 1970, pp. 509-522.
- 163. Willett, H. C.: The Pattern of Solar Climatic Relationships. Ann. N. Y. Acad. Sci., Vol. 95, 1961, pp. 89-106.
- 164. King, J. W.: Sun-Weather Relationships. Astron. Aeron., Vol. 13, April 1975, pp. 10-19.
- 165. Link, F.: Auroral and Climatic Cycles in the Past. J. Brit. Astron. Ass., Vol. 78, 1968, pp. 195-205.

- 166. Matsushita, S.: Ancient Aurorae Seen in Japan. J. Geophys. Res., Vol. 61, 1956, pp. 297-302.
- 167. Bandeen, W. R. and Maran, S. P. (eds.): Possible Relationships Between Solar Activity and Meteorological Phenomena. NASA SP-366, Washington, D.C., 1975.
- 168. Downie, N. M. and Heath, R. W.: <u>Basic Statistical Methods</u> (2nd Edition). Harper and Row, Publ., New York, New York, 1965.
- 169. Longley-Cook, L. H.: <u>Statistical Problems</u>. College Outline Series, Barnes and Noble Books, A Division of Harper and Row, Publ., New York, New York, 1970.
- 170. Waldmeier, W.: The Sunspot-Activity in the Years 1610-1960. Schulthess and Co., Zurich, Switzerland, 1961.
- 171. Sargent, H. H. III: Reference Points in the Sunspot Cycle. <u>Bull. Am. Astron.</u> <u>Soc.</u>, Vol. 16, 1984, p. 452.
- 172. Sonett, C. P.: Sunspot Time Series: Spectrum from Square Law Modulation of the Hale Cycle. Geophys. Res. Letters, Vol. 9, 1982, pp. 1313-1316.
- 173. Gilliland, R. L.: Solar Radius Variations over the Past 265 Years. Astrophys. J., Vol. 248, 1981, pp. 1144-1155.
- 174. Williams, G. E.: Sunspot Periods in the Late Precambrian Glacial Climate and Solar-Planetary Relations. Nature, Vol. 291, 1981, pp. 624-628.
- 175. Long, A.: 100 to 200 Year Solar Periodicities. <u>Nature</u>, Vol. 298, 1982, pp. 223-224.
- 176. Wigley, T. M. L.: Sun-Climate Links. Nature, Vol. 288, 1980, pp. 317-319.
- 177. Silverman, S. M.: The Visual Aurora as a Predictor of Solar Activity. <u>J</u>. Geophys. Res., Vol. 88, 1983, pp. 8123-8128.
- 178. Bhalme, H. N. and Jadhav, S. K.: The Double (Hale) Sunspot Cycle and Floods and Droughts in India. Weather, Vol. 39, 1984, pp. 112-116.
- 179. Voichishin, K. S.: Peculiarities of the Fine Structure of the 11-Year Cyclicity of Solar Activity. Sov. Astron., Vol. 25, 1981, pp. 93-96.
- Poulakos, C. and Tritakis, B.: A Study of the Areas Index I. Solar Activity and Related Interplanetary and Terrestrial Phenomena (ed., J. Xanthakis), Springer-Verlag, New York, New York, 1973, pp. 49-60.
- 181. Papagiannis, M. D., Zerefos, C. S. and Reparis, C. S.: Evidence of a Double Peak in Solar Maxima. Solar Activity and Related Interplanetary and Terrestrial Phenomena (ed., J. Xanthakis), Springer-Verlag, New York, New York, 1973, pp. 75-82.

- Bonov, A. D.: Characteristic Features of Secular and Supersecular Cycles of Solar Activity. A 180-Year Cycle. Solar Activity and Related Interplanetary and Terrestrial Phenomena (ed., J. Xanthakis), Springer-Verlag, New York, New York, 1973, pp. 83-86.
- 183. Xanthakis, J.: The Different Indices of Solar Activity and the Time of Rise.

 Solar Physics (ed., J. Xanthakis), Interscience Publ., John Wiley and Sons,

 New York, New York, 1967, pp. 157-227.

APPROVAL

ON LONG-TERM PERIODICITIES IN THE SUNSPOT RECORD

By Robert M. Wilson

The information in this report has been reviewed for technical content. Review of any information concerning Department of Defense or nuclear energy activities or programs has been made by the MSFC Security Classification Officer. This report, in its entirety, has been determined to be unclassified.

A. J. DESSLER

Director, Space Sciences Laboratory